


Welcome to **TEZSPIRE**TM (tezepelumab injection)

This brochure is intended for people who have been prescribed TEZSPIRETM. This brochure does not replace the advice of your healthcare team. If you have questions or concerns, be sure to contact your healthcare team.



What is TEZSPIRE™ and how does it work?

TEZSPIRE™ is a prescription medicine used with other asthma medicines for the maintenance treatment of severe asthma in people 12 years of age and older whose asthma is not controlled with their current asthma medicine. TEZSPIRE™ is not used to treat sudden breathing problems.

- ▶ TEZSPIRE™ works by blocking the action of thymic stromal lymphopoietin (TSLP).
- ▶ TSLP is a protein that plays a role in causing the signs and symptoms of asthma.
- ▶ TEZSPIRE™ may reduce the number of asthma attacks you experience, improve your breathing and reduce your asthma symptoms.



How do I take TEZSPIRE™?



Always use TEZSPIRE™ exactly as your healthcare team has described. If you or your caregiver have any questions, talk to your healthcare team.



TEZSPIRE™ is given as an injection just under the skin (subcutaneously) either by a single-use pre-filled pen or a single-use pre-filled syringe. You and your healthcare team—including your Connect360 healthcare professional—will have decided together which injection device is the best for you.



You and your healthcare team will decide if you or your caregiver should inject TEZSPIRE™. Before using a TEZSPIRE™ pre-filled pen or pre-filled syringe, your healthcare team will show you or your caregiver how to use it the right way. Read the Instructions for Use (that came in your TEZSPIRE™ product box) for the pre-filled pen or pre-filled syringe carefully before using TEZSPIRE™ and each time you get a refill.



Do not share or use a TEZSPIRE™ pre-filled pen or pre-filled syringe more than one time.



Do not stop using TEZSPIRE™ without speaking to your healthcare team first. Interrupting or stopping the treatment with TEZSPIRE™ may cause your asthma symptoms and attacks to come back.

TEZSPIRE™ is an injectable medication that you or a caregiver can administer after training.

Resources are available to help patients use the pre-filled pen or pre-filled syringe correctly:

- ▶ Read the Instructions for Use that came in your TEZSPIRE™ product box,
- ▶ Review the Patient Journey Guide that came in your Connect360 Starter Kit, and
- ▶ Scan the appropriate QR code below to watch a video on how to inject TEZSPIRE™.





**Using your TEZSPIRE™
Pre-filled Pen**



**Using your TEZSPIRE™
Pre-filled Syringe**

How do I take TEZSPIRE™?

	MON	TUES	WED	THURS	FRI	SAT	SUN
1							
2							
3							
4							
5							

The usual recommended dose of TEZSPIRE™ is **210 mg every 4 weeks** delivered by pre-filled pen or pre-filled syringe. Each pre-filled pen or pre-filled syringe contains one dose of TEZSPIRE™.

What if I miss a dose of TEZSPIRE™?

If you or your caregiver missed a dose:	Inject a dose as soon as possible and continue your next injection on your usual injection day.
If you did not notice that you have missed a dose until it is time for your next scheduled dose:	Inject the next scheduled dose as planned.
If you are not sure when to inject TEZSPIRE™:	Contact your healthcare team.

What if I think I've taken too much TEZSPIRE™?

If you, or your caregiver, think you have taken too much TEZSPIRE™, contact your healthcare team, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

How do I store TEZSPIRE™ before injecting at home?

Store TEZSPIRE™ in a refrigerator between 2°C and 8°C in its original package to protect it from light. TEZSPIRE™ may alternatively be kept at room temperature between 20°C and 25°C in the original package for a maximum of 30 days. When TEZSPIRE™ has reached room temperature, do not put it back in the refrigerator. Dispose of TEZSPIRE™ that has been stored at room temperature for more than 30 days.

DO NOT:



Use TEZSPIRE™ if it has been dropped, damaged, or if the security seal on the carton has been broken.



Use TEZSPIRE™ past the expiry date which is stated on the label. The expiry date refers to the last day of that month.



Shake, freeze or expose TEZSPIRE™ to heat.

DO:



Dispose of unused medicinal product or waste material in accordance with local requirements.



Keep out of reach and sight of children.

What are the potential side effects of TEZSPIRE™?

Common side effects of TEZSPIRE™ include:



Sore throat



Injection site reaction
(i.e., redness, swelling, and pain)



Joint pain



Rash

These are not all the potential side effects you may have when taking TEZSPIRE™. If you experience any side effects not listed here, tell your health professional.



If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare team.

What can I do to help avoid side effects and ensure proper use?

Before taking TEZSPIRE™, tell your healthcare provider about any health conditions or problems you may have, including if you:



Have had any symptoms of an allergic reaction (symptoms may vary, but can include breathing problems, hives, and rash). If you notice any of these signs, speak to your healthcare team immediately.



Have a parasitic infection or if you live in an area where parasitic infections are common or if you are travelling to such a region. TEZSPIRE™ may weaken your ability to fight certain types of parasitic infections.



Are pregnant, think you may be pregnant, or plan to become pregnant. Do not use TEZSPIRE™ unless your healthcare team tells you to. It is not known if TEZSPIRE™ can harm your unborn baby.



Are breast-feeding. TEZSPIRE™ may pass into breast milk. Your healthcare team will discuss with you whether you should stop treatment with TEZSPIRE™ while you are breast-feeding, or if you should stop breast-feeding.

Do not use TEZSPIRE™ if you are allergic to tezepelumab or any of its ingredients.

You should also talk to your healthcare team if you have asthma that remains uncontrolled or worsens during treatment with TEZSPIRE™.

Other warnings you should know about:

- ▶ TEZSPIRE™ is not a rescue medicine and should not be used to treat a sudden asthma attack.
- ▶ TEZSPIRE™ is not approved for use in children below the age of 12.

Tell your healthcare team about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

- ▶ There are no known interactions with TEZSPIRE™.
- ▶ Tell your healthcare team if you have recently had or are due to have a vaccination.
- ▶ Do not suddenly stop taking your asthma medicines once you have started TEZSPIRE™.
 - ▶ These medicines (especially ones called corticosteroids) must be stopped gradually, under the direct supervision of your doctor and based on your response to TEZSPIRE™.



Notes

The Connect360 Patient Support Program



The Connect360 Patient Support Program provides resources designed for your support and to help you manage your severe asthma. Alongside your doctor's treatment plan, you can have access to:



Reimbursement Navigation

- ▶ Investigation of your insurance and coverage options.
- ▶ Assistance to coordinate insurance benefits to help facilitate continued access to your medication.



Injection and Self-Administration Training

- ▶ Coordination of your injection appointments and complete training on how to self-administer TEZSPIRE™.
- ▶ Regular appointment reminders.
- ▶ Post-injection reports for your doctor.



Ongoing Support

- ▶ Program Care Coordinator to answer any questions about TEZSPIRE™.
- ▶ Regular check-in calls throughout your treatment.
- ▶ Access to a toll-free helpline, **1 (833) 360-2666**, available Mon–Fri, 8am–8pm EST.
- ▶ Access to email support through **connect360@innomar-strategies.com**.



Educational Resources

- ▶ Disease and treatment education to help you manage your severe asthma.

Connect360 is for patients prescribed with TEZSPIRE™ by their doctor. It does not replace your doctor's treatment plan. You can choose to adapt the service or opt out as your needs change.



Questions?



Call the Connect360 Patient Support Program at
1 (833) 360-2666 Monday to Friday, 8 am to 8 pm (EST) or



Email **connect360@innomar-strategies.com**

TEZSPIRE™ and the logo are trademarks of Amgen Inc. and the Amgen logo is a registered trademark of Amgen Inc., used under license by AstraZeneca Canada Inc. The AstraZeneca logo is a trademark of AstraZeneca AB, used under license by AstraZeneca Canada Inc.

©2023 AstraZeneca. All rights reserved. Last Updated 10/23



TEZ2210E